

Landrace. 1202; mani blanco. Collected 04/21/1992 in Bolivia. Latitude 15 deg. 47' S. Longitude 66 deg. 58' W. Elevation 560 m. Covendo, Sud Yungas Province, La Paz Dept. Fruits 4-5.5cm long, 2cm diam., straight or curved, with little or no constriction, with humps, keel and beak. Reticulation rounded to almost smooth, containing 3, 4 white seeds.

The following were collected by Samuel Sanchez-Dominguez, Universidad Autonoma Chapingo, Departamento de Fitotecnia, Chapingo 56230, Mexico, Mexico ; David E. Williams, USDA, ARS, Natl. Germplasm Resources Laboratory, Building 003, Room 400, BARC-West, Beltsville, Maryland 20705-2350, United States. Donated by David E. Williams, USDA, ARS, Natl. Germplasm Resources Laboratory, Building 003, Room 400, BARC-West, Beltsville, Maryland 20705-2350, United States. Received 12/29/1992.

**PI 576608. *Arachis hypogaea* L. var. *fastigiata***

Landrace. 1223; cacahuete chocolin. Collected 11/07/1992 in Mexico. Latitude 18 deg. 40' N. Longitude 98 deg. 32' W. Elevation 1450 m. Tlapanala, Edo. de Puebla. Plant erect. Pods 4.5-6.0cm long, fairly straight. Constriction slight to moderate. Humps pronounced. Beak and reticulation moderate. 3-4 red seeds. Matures in 4 1/2 months. Non-commercial variety planted for home consumption, usually boiled. Said to have been grown here for about 20 years.

**PI 576609. *Arachis hypogaea* L. var. *fastigiata***

Landrace. 1224; cacahuete colorado. Collected 11/07/1992 in Mexico. Latitude 18 deg. 38' N. Longitude 98 deg. 41' W. Elevation 1400 m. Edo. de Puebla, Mpio. Tepexco, Tepexco. Plant erect with light green leaves. Pods 4.0-6.0cm long, straight. Constriction slight. Humps, beak and reticulation prominent. 4 red seeds. Most widely cultivated variety in this town.

**PI 576610. *Arachis hypogaea* L. var. *fastigiata***

Landrace. 1231; cacahuete chino. Collected 11/08/1992 in Mexico. Latitude 18 deg. 45' N. Longitude 98 deg. 46' W. Elevation 1600 m. Edo. de Morelos, Mpio. Temoac, Huazulco. Pods 4.5-7.0cm long, fairly straight. Constriction slight to moderate. Humps and beak pronounced. Reticulation prominent but not sharp. 4 red seeds.

**PI 576611. *Arachis hypogaea* L. var. *fastigiata***

Landrace. 1233; cacahuete colorado. Collected 11/08/1992 in Mexico. Latitude 18 deg. 45' N. Longitude 98 deg. 46' W. Elevation 1600 m. Edo. de Morelos, Mpio. Temoac, Huazulco. Plant said to be erect. 3-4 red seeds. Said that this variety was brought here years ago from Tepexco, Puebla.

**PI 576612. *Arachis hypogaea* var. *hirsuta* Kohler**

Landrace. 1209; cacahuete criollo. Collected 11/05/1992 in Mexico. Latitude 18 deg. 45' N. Longitude 98 deg. 30' W. Elevation 1680 m. Cacaloxuchil, Mpio. Huaquechula, Edo. de Puebla. Plant said to be runner type. Pods 3.5-4.5cm long, fairly straight, with little or no constriction. Humps and beak slight. Reticulation pronounced but not sharp, containing 3 dark purple seeds. Common.

**PI 576613. *Arachis hypogaea* var. *hirsuta* Kohler**

Landrace. 1210; cacahuete criollo morado. Collected 11/05/1992 in Mexico. Latitude 18 deg. 45' N. Longitude 98 deg. 30' W. Elevation 1670 m. Edo. de Puebla, Mpio. Huaquechula, Santiago Tetla. Plant prostrate. Pods 3.0-4.5cm long, straight, little or no constriction. Humps slight. Beak small. Reticulation deep and sharp. 3 purple seeds. Said to be the original local variety cultivated here since the time of the "abuelos." Matures in 5 months.

**PI 576614. *Arachis hypogaea* var. *hirsuta* Kohler**